

**GENERIC PHARMACEUTICAL SERVICES, INC.**  
**DRUG FACTS LABEL GUIDE COPY**

Product: ACETAMINOPHEN 500 mg TABLETS

Dist. by: *Name, city, state, zip code*      *2 tablets per packet*

<b>DRUG FACTS</b>	
<b>Active Ingredient (in each tablet)</b> Acetaminophen 500 mg.....	<b>Purpose</b> Pain Reliever/Fever Reducer
<b>Uses</b> temporary relief of minor aches and pains due to: • headache backache common cold menstrual cramps • muscular aches arthritis toothache reduction of fever	
<b>Warnings</b> <b>Allergy Alert:</b> Do not take this product if you are hypersensitive to acetaminophen. <b>Alcohol Warning:</b> If you generally consume 3 or more alcohol drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage.	
<b>Do not use</b> • with any other product containing acetaminophen	
<b>Stop use and ask a doctor if</b> • new symptoms occur • redness or swelling is present • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. <b>Overdose warning:</b> Taking more than the recommended dose (overdose) may cause liver damage. In the case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.	
<b>Directions</b> adults and children 12 years and over	• take 2 tablets every 4 to 6 hours as needed • do not take more than 8 tablets in 24 hours
children under 12 years	Do not use this Adult Extra Strength product in children under 12 years of age; this will provide more than the recommended dose (overdose) of Pain Reliever and could cause serious problems.
<b>Other Information</b> store at room temperature, 15°-30° C (59-86° F)	
<b>Inactive Ingredients:</b> may contain: corn starch, pregelatinized starch, povidone, polyvinylpyrrolidone, sodium carboxymethylcellulose, sodium starch glycolate, stearic acid	

**TAMPER EVIDENT PACKETS. DO NOT USE IF OPEN OR TORN.**

*Lot # xxxxx, Exp. xx/xx*

*Rev. 3/07*

**By:** mfM      **Date:** 3/20/07

**Check By:** h      **Date:** 3/22/07

**Manufacturer: CPC # 0546 and A&Z Pharmaceutical**

**Summary Revisions:**

Revision: (3/07) Added supplier CPC inactives. No "Other" declaration required for CPC or A&Z.

Revision: (6/06) Added secondary supplier A&Z Pharmaceutical inactives.

Revision: (1/06) Added Overdose Warning (like Tylenol)

Revision: (6/05)

- Uses: added backache, the common cold, menstrual cramps, arthritis, toothache
- Warnings: added "Do not use with any other product containing acetaminophen" (deleted 2 statements not on PDK or McNeil Tylenol®)
- Precautions: added new symptoms occur, "Quick medical attention"
- Directions: # tabs every 4 to 6 hours and Warning for children under 6 years
- Inactives changed
- No declaration required-Final rule: 9/24/05 OTC labeling containing calcium, magnesium, potassium and sodium (actives and inactives).